

K3 SURFACES ASSOCIATED TO CURVES OF GENUS TWO

ABHINAV KUMAR

ABSTRACT. It is known ([GLD], [Na]) that there is a unique K3 surface X which corresponds to a genus 2 curve C such that X has a Shioda-Inose structure with quotient birational to the Kummer surface of the Jacobian of C . In this paper we give an explicit realization of X as an elliptic surface over $\mathbb{P}^1$ with specified singular fibers of type II^* and III^* . We describe how the Weierstrass coefficients are related to the Igusa-Clebsch invariants of C .

CONTENTS

1. Introduction	1
2. Preliminaries on lattices	3
3. K3 surfaces	5
3.1. Background	5
3.2. Kummer surfaces	7
3.3. Shioda-Inose structures	8
3.4. Elliptic K3 surfaces	9
4. Genus 2 curves	11
4.1. Background: Moduli and invariants	11
4.2. Kummer surface	11
5. Family of elliptic K3 surfaces associated to genus 2 curves	14
5.1. Elliptic surfaces with E_8 and E_7 fibers	14
5.2. Main theorem	15
5.3. Néron-Severi lattices	16
5.4. Completion of proof	18
6. An application to the moduli of six points in $\mathbb{P}^1$	20
Acknowledgements	21
References	21

1. INTRODUCTION

In this paper, we study some K3 surfaces of high rank. Precisely, we consider K3 surfaces which have a Shioda-Inose structure: that is, an involution which preserves a global 2-form, such that the quotient is a Kummer surface. Kummer surfaces are

Date: January 23, 2007.

This work was the outcome of the author's doctoral thesis at Harvard University, and was supported by a Putnam Fellowship.

a special class of K3 surfaces which are quotients of abelian surfaces. The Kummer surface thus carries algebro-geometric information about the abelian surface. It has Néron-Severi rank at least 17, and so therefore do the K3 surfaces with Shioda-Inose structure.

These surfaces were first studied by Shioda and Inose [IS], who give a description of *singular* K3 surfaces, i.e. those with rank 20, the maximum possible for a K3 surface over a field of characteristic zero. They prove that there is a natural one-to-one correspondence between the set of singular K3 surfaces up to isomorphism and the set of equivalence classes of positive definite even integral binary quadratic forms. The result follows that of Shioda and Mitani [MS] who show that the set of singular abelian surfaces (that is, those having Picard number 4) is also in one-to-one correspondence with the equivalence classes of positive definite even integral binary quadratic forms. The construction of Shioda and Inose produces a singular K3 surface by taking a double cover of a Kummer surface associated to a singular abelian surface and with a specific type of elliptic fibration. The resulting K3 surface has an involution such that the quotient is the original Kummer surface. It also turns out that the lattice of transcendental cycles on the K3 surface (i.e. the orthogonal complement of the Néron-Severi group in the second singular cohomology group) is isomorphic to the lattice of transcendental cycles on the abelian surface.

Nikulin studied groups of automorphisms of K3 surfaces in [N1], and Morrison studied Shioda-Inose structures more extensively in [M1], giving necessary and sufficient conditions for a K3 surface to have Shioda-Inose structure, in terms of the Néron-Severi group of the K3 surface.

The Shioda-Inose setup takes a K3 surface and produces the data of a Kummer surface, or of an abelian surface. One may reverse the question, and ask: given an abelian surface A , can we produce a K3 surface with Shioda-Inose structure such that its quotient under the Nikulin involution is birational to $\mathrm{Km}(A)$?

In particular, when A is the Jacobian of a curve C of genus 2, Galluzzi and Lombardo [GLD] show that there is a unique K3 surface (up to isomorphism) with Shioda-Inose structure which corresponds to A under this construction. Dolgachev, in the appendix to the paper, proves an isomorphism between the moduli space of elliptic K3 surfaces with bad fibers of type E_8 and E_7 , and the moduli space of principally polarized abelian surfaces. However, the isomorphism is not explicit, and uses a result of Naruki [Na] to show that the quotient surface constructed is a Kummer surface.

In this paper we describe explicitly the relation between C and the K3 surface X with E_8 and E_7 fibers. It turns out that we can give an alternative elliptic fibration on X , with a 2-torsion section. The Nikulin involution on X is translation by the 2-torsion section. The quotient elliptic surface under the 2-isogeny is the Kummer surface of a unique principally polarized abelian surface (generically, the Jacobian of a curve of genus 2). We relate the Igusa-Clebsch invariants of the genus 2 curve

to the moduli of the original K3 elliptic surface, thus giving an explicit description of the map on moduli spaces.

This work generalizes a construction of Elkies [E], who found a family of K3 elliptic surface with two E_8 fibers such that the quotient is birational to the Kummer surface of a product of two elliptic curves E_1 and E_2 , and is related to it by a Galois-invariant map of degree 9. Earlier, Inose [In] had found the isogenous K3 surface to the Kummer surface of $E_1 \times E_2$, but with maps that were defined over an algebraically closed ground field. Clingher and Doran have also studied the K3 surfaces corresponding to $E_1 \times E_2$ in [CD], as well as to $J(C)$ more recently.

Such explicit formulas have some Diophantine applications. Elkies has used his techniques from [E] to construct elliptic curves of high rank over $\mathbb{Q}(t)$ and $\mathbb{Q}$. Similarly, he used a K3 surface of rank 19 corresponding to a rational point on a Shimura curve to find an elliptic curve of rank at least 28 over $\mathbb{Q}$. As an application of the results of this paper, we can find explicit parameterizations of some Hilbert modular surfaces. We hope to address these techniques and results in a future publication.

The outline of the paper is as follows: in Section 2, we recall facts about integral lattices which appear in this context as sublattices of Néron-Severi groups or cohomology groups of K3 surfaces. In Section 3, we cover the necessary background on K3 surfaces, whereas section 4 recalls the theory of genus 2 curves. In Section 5, we state and prove the main theorem, and in section 6, we describe a connection to the moduli space of 6 points in $\mathbb{P}^1$.

2. PRELIMINARIES ON LATTICES

Definition 1. A **lattice** will denote a finitely generated free abelian group Λ equipped with a symmetric bilinear form $B : \Lambda \times \Lambda \rightarrow \mathbb{R}$. An **integral lattice** is a lattice whose form takes values in $\mathbb{Z}$.

We abbreviate the data (Λ, B) to Λ sometimes, when the form is understood, and we interchangeably write $u \cdot v = \langle u, v \rangle = B(u, v)$ and $u^2 = u \cdot u = \langle u, u \rangle = B(u, u)$ for $u, v \in \Lambda$.

The **signature** of the lattice is the real signature of the form B , written (r_+, r_-, r_0) where r_+ , r_- and r_0 are the number of positive, negative and zero eigenvalues of B , counted with multiplicity. We say that the lattice is **non-degenerate** if the form B has zero kernel, i.e. $r_0 = 0$. In that case, the signature is abbreviated to (r_+, r_-) . We say Λ is **even** if $u^2 \in 2\mathbb{Z}$ for all $u \in \Lambda$. The dual lattice to Λ is denoted Λ^* .

Let Λ be a non-degenerate lattice. The **discriminant** of the lattice is $|\det(B)| = |\Lambda^*/\Lambda|$. The lattice is said to be **unimodular** if its discriminant is 1. Note that in

the literature, the discriminant is frequently defined to be $\det(B) = (-1)^{r-1} |\Lambda^*/\Lambda|$, in which case a lattice is unimodular iff its discriminant is ± 1 .

For a lattice Λ and a real number α , we denote by $\Lambda(\alpha)$ the lattice which has the same underlying group but with the bilinear form scaled by α . The lattice of rank one with a generator of norm α will be denoted $\langle \alpha \rangle$.

By a **root** of a positive definite lattice, we will mean an element x such that $x^2 = 2$, whereas for a negative-definite or indefinite lattice, we will mean an element x such that $x^2 = -2$.


We note here some theorems about the structure and embeddings of lattices.

A **root lattice** is a lattice that is generated (as an abelian group) by its roots. First, let us introduce some familiar root lattices, through their Dynkin diagrams. The subscript in the name of the lattice is the dimension of the lattice, which is also the number of nodes in the Dynkin diagram.




$A_n (n \geq 1)$, signature $(n, 0)$, discriminant $n + 1$.

This is the positive definite lattice with n generators $v_1, \dots, v_n$ with $v_i^2 = 2$ and $v_i \cdot v_j = -1$ if the vertices i and j are connected by an edge, and 0 otherwise. It may be realized as the set of integral points on the hyperplane $\{x \in \mathbb{R}^{n+1} \mid \sum x_i = 0\}$.




$D_n (n \geq 4)$, signature $(n, 0)$, discriminant 4.

D_n can be realized as $\{x \in \mathbb{Z}^n \mid \sum_{i=1}^n x_i \equiv 0 \pmod{2}\}$. It has $2n(n-1)$ roots.




E_8 , signature $(8, 0)$, discriminant 1.

One realization of E_8 is as the span of D_8 and the all-halves vector $(1/2, \dots, 1/2)$. It has 240 roots.



E_7 , signature $(7, 0)$, discriminant 2.

Taking the orthogonal complement of any root in E_8 gives us E_7 . It has 126 roots.



E_6 , signature $(6, 0)$, discriminant 3.

Taking the orthogonal complement of e_1 and e_2 in E_8 , where e_1, e_2 are roots such that $e_1 \cdot e_2 = -1$, gives us E_6 . It has 72 roots.

We let the **Nikulin lattice** N be the lattice generated by $v_1, \dots, v_8$ and $\frac{1}{2}(v_1 + \dots + v_8)$, with $v_i^2 = -2$ and $v_i \cdot v_j = 0$ for $i \neq j$. It is isomorphic to $D_8^*(-2)$. It has signature $(0, 8)$ and discriminant 2^6 .

One checks easily that N has 16 roots, namely $\pm v_i$. In particular, N is not a root lattice.

Let U be the hyperbolic plane, i.e. the indefinite rank 2 lattice whose matrix is

$$\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$$

Note that $U(-1) \cong U$.

Indefinite lattices are sometimes easier to characterize than definite ones. For instance, one has the following theorem of Milnor.

Theorem 2. (Milnor [Mi]) *Let Λ be an indefinite unimodular lattice. If Λ is even, then $\Lambda \cong E_8(\pm 1)^m \oplus U^n$ for some m and n . If Λ is odd, then $\Lambda \cong \langle 1 \rangle^m \oplus \langle -1 \rangle^n$ for some m and n .*

However, computationally, indefinite lattices are difficult to work with. For instance, it is typically much easier to find roots of a definite lattice than of an indefinite one.

3. K3 SURFACES

3.1. Background. In this section we recall the definition and basic properties of K3 surfaces. For more details we refer the reader to [BHPV], [K] and [M2]. Let X be a smooth projective surface over a field k .

Definition 3. We say that X is a **K3 surface** if $H^1(X, \mathcal{O}_X) = 0$ and the canonical bundle of X is trivial, i.e. $K_X \cong \mathcal{O}_X$.

In the sequel, we will assume that k is the field $\mathbb{C}$ of complex numbers or some subfield of $\mathbb{C}$.

K3 surfaces are simply connected. From Noether's formula

$$(3.1) \quad \chi(\mathcal{O}_X) = \frac{1}{12}(K^2 + c_2)$$

it follows that the Euler characteristic of X is 24. Here $c_1 = 0$ and $c_2 = \chi(X)$ are the Chern classes of (the tangent bundle) of X . Therefore the middle cohomology $H_X := H^2(X, \mathbb{Z})$ is 22-dimensional. Identifying $H^4(X, \mathbb{Z})$ with $\mathbb{Z}$, we also know that H_X endowed with the cup-product pairing is a unimodular lattice (that is, the associated quadratic form has discriminant 1) because of Poincaré duality. Finally, we also know that H_X is an even lattice; this follows from Wu's formula involving Stiefel-Whitney classes. Finally, using the index theorem

$$(3.2) \quad \tau(X) = \frac{1}{3}(c_1^2 - 2c_2)$$

we see that H_X has signature $(3, 19)$.

Using the classification of Milnor, Theorem 2, we use these properties to conclude that $H_X \cong E_8(-1)^2 \oplus U^3$ as lattices.

Now, the exact sequence of analytic sheaves

$$0 \rightarrow \mathbb{Z} \rightarrow \mathcal{O}_X \xrightarrow{z \mapsto e^{2\pi iz}} \mathcal{O}_X^* \rightarrow 0$$

gives, upon taking the long exact sequence of cohomology, an injective map (the first Chern class map)

$$H^1(X, \mathcal{O}_X^*) \rightarrow H^2(X, \mathbb{Z}).$$

Thus, an analytic line bundle is determined by its image in $H^2(X, \mathbb{Z})$. For an algebraic K3 surface, we can also say that an algebraic line bundle is determined by class in $H^2(X, \mathbb{Z})$. Linear equivalence, algebraic equivalence, and numerical equivalence all coincide for an algebraic K3 surface.

Definition 4. The image of $H^1(X, \mathcal{O}_X^*)$ in $H^2(X, \mathbb{Z}) \subset H^2(X, \mathbb{R})$ is a sublattice of $H^2(X, \mathbb{Z})$, which we call the **Néron-Severi** group of X , and denote by $NS(X)$ or S_X or $\text{Pic}(X)$.

K3 surfaces are Kähler, and we have a Hodge structure

$$H^2(X, \mathbb{Z}) \otimes \mathbb{C} \cong H^{0,2}(X) \oplus H^{1,1}(X) \oplus H^{2,0}(X)$$

In fact, it can be shown that

$$NS(X) = H_{\mathbb{Z}}^{1,1}(X) := H^{1,1}(X) \cap H^2(X, \mathbb{Z}).$$

Also,

$$NS(X) = \{z \in H^2(X, \mathbb{Z}) \mid \langle z, H^{2,0}(X) \rangle = 0\}.$$

Since the canonical bundle of X is trivial, there exists a regular $(2, 0)$ form on X , and we have $h^{2,0} = 1$. It follows that the Hodge diamond of X is

$$\begin{array}{ccccc} & & 1 & & \\ & 0 & & 0 & \\ 1 & & 20 & & 1 \\ & 0 & & 0 & \\ & & 1 & & \end{array}$$

We will only concern ourselves with algebraic K3 surfaces, i.e. those which are nonsingular projective algebraic varieties over k .

Example 5. A smooth complete quartic hypersurface in $\mathbb{P}^3$ is a K3 surface.

Example 6. A smooth complete intersection of three quadric hypersurfaces in $\mathbb{P}^5$ is a K3 surface.

Definition 7. A (-2) -**curve** on a surface is an irreducible rational curve with self-intersection -2 .

On a K3 surface, it is an easy exercise using the genus formula to see that curves of self-intersection -2 are exactly the smooth rational curves.

3.2. Kummer surfaces. Here we recall the construction of the Kummer surface associated to an abelian surface, which is an example of a K3 surface.

Let A be an abelian surface, and let ι be the involution which is multiplication by -1 . Note that ι fixes any regular algebraic 2-form on A (the space of 2-forms is one-dimensional since $h^0(X, K) = h^2(X, \mathcal{O}) = 1$). Now if we blow up the surface A at its sixteen 2-torsion points, we get a surface $\tilde{A}$ on which ι extends to an involution $\tilde{\iota}$. The quotient surface $Y = \tilde{A}/\{1, \tilde{\iota}\}$ is in fact a K3 surface, and the regular 2-form is induced from the one on A . The surface $Y = \text{Km}(A)$ is called the Kummer surface of A . It has sixteen disjoint rational curves $F_1, \dots, F_{16}$ which are the images of the sixteen exceptional divisors on $\tilde{A}$.

The Néron-Severi lattice of a Kummer surface has 16 linearly independent divisor classes coming from the sixteen rational curves above. These generate a negative definite lattice, and there is also a class of a polarization on $\text{Km}(A)$, since it is projective. Therefore its signature is $(1, r)$ for some $r \geq 16$. In fact, the Néron-Severi lattice always contains a particular lattice of signature $(0, 16)$ and discriminant 2^6 , called the Kummer lattice K . We describe its structure.

The set $I := A[2] \cong (\mathbb{Z}/2)^4 \cong \mathbb{F}_2^4$ of 16 elements has a natural structure of a vector space of dimension 4 over $\mathbb{F}_2$. Choose a labeling $I = \{1, 2, \dots, 16\}$ (for instance, by writing $i - 1 = \sum_{j=0}^3 b_{ij} 2^j$ for $1 \leq i \leq 15$) and let $f_1, \dots, f_{16}$ be the classes of the rational curves corresponding to the blowups at the 2-torsion points. Let Q be the set of 32 elements consisting of 30 affine hyperplanes (considered as subsets of I) as well as the empty set and all of I . This set has the structure of a vector

space over $\mathbb{F}_2$, the addition operation being symmetric difference of sets. The set Q is in fact the Reed-Muller code $\mathcal{R}(1, 4)$: the characteristic functions of the sets in Q , viewed as functions from $\mathbb{F}_2^4$ to $\mathbb{F}_2$, are exactly the polynomials in 4 variables of degree at most 1. For every $M \in Q$, we have an element $f_M = \frac{1}{2} \sum_{i \in M} f_i$ of $\sum \mathbb{Q} f_i \in NS(\text{Km}(A)) \otimes \mathbb{Q}$. These vectors actually lie in $NS(\text{Km}(A))$. The lattice generated by the $f_i, (i = 1, \dots, 16)$ and the f_M has discriminant $2^{16}/(2^5)^2 = 2^6$ and it is called the Kummer lattice.

In section 4, we will say more about the Kummer surface associated to a principally polarized abelian surface which arises as the Jacobian of a genus 2 curve.

3.3. Shioda-Inose structures. We now describe the data of a Shioda-Inose structure on a K3 surface.

Definition 8. An involution ι on a K3 surface X is called a **Nikulin involution** if $\iota^*(\omega) = \omega$ for every $\omega \in H^{2,0}(X)$.

In fact, a Nikulin involution fixes T_X pointwise. Every Nikulin involution has 8 isolated fixed points. As in the construction of the Kummer surface, we may blow up these points to get $\tilde{X}$, which has eight exceptional curves, and an involution $\tilde{\iota}$. The quotient of $\tilde{X}/\tilde{\iota}$ by a Nikulin involution is a K3 surface Y .

The images of the exceptional curves are (-2) -curves on the quotient K3 surface Y . The Néron-Severi lattice of Y contains the Nikulin lattice N : it is generated by vectors $c_1, \dots, c_8$ (the classes of the eight disjoint -2 -curves) and $\frac{1}{2} \sum c_i$, with the form induced by $c_i \cdot c_j = -2\delta_{ij}$.

Definition 9. We say that X admits a **Shioda-Inose structure** if there is a Nikulin involution ι on X with rational quotient map $\pi : X \rightarrow Y$ such that Y is a Kummer surface and π_* induces a Hodge isometry $T_X(2) \cong T_Y$.

If X has a Shioda-Inose structure, let A be the abelian surface whose Kummer surface is Y . Then we have a diagram

$$\begin{array}{ccc} X & & A \\ & \searrow \quad \swarrow & \\ & Y & \end{array}$$

of rational maps of degree 2, and Hodge isometries $T_X(2) \cong T_Y \cong T_A(2)$, thus inducing a Hodge isometry $T_X \cong T_A$.

The following theorem of Morrison characterizes K3 surfaces with Shioda-Inose structures in terms of the Néron-Severi lattice.

Theorem 10. (Morrison [M1]) *Let X be an algebraic K3 surface. The following are equivalent:*

- (1) X admits a Shioda-Inose structure.

- (2) *There exists an abelian surface A and a Hodge isometry $T_X \cong T_A$.*
- (3) *There is a primitive embedding $T_X \hookrightarrow U^3$.*
- (4) *There is an embedding $E_8(-1)^2 \hookrightarrow NS(X)$.*

3.4. Elliptic K3 surfaces. We recall here a few facts about elliptic surfaces. References for these are [S] and [Si].

Definition 11. An elliptic surface is a smooth projective algebraic surface X with a proper morphism $\pi : X \rightarrow C$ to a smooth projective algebraic curve C , such that

- (1) There exists a section $\sigma : C \rightarrow X$.
- (2) The generic fiber E is an elliptic curve.
- (3) π is relatively minimal.


Concretely, we will be considering the case $C = \mathbb{P}^1$, since a K3 surface cannot have a non-constant map to a curve of positive genus. We will choose a Weierstrass equation for the generic fiber, which is an elliptic curve over the function field $\mathbb{C}(\mathbb{P}^1) = \mathbb{C}(t)$, namely

$$y^2 + a_1(t)xy + a_3(t)y = x^3 + a_2(t)x^2 + a_4(t)x + a_6(t)$$

where a_i are rational functions of t . In fact, by multiplying x and y by suitable rational functions, we can make $a_i(t)$ polynomials in t . Furthermore, we may translate x by a rational function, and y by a $\mathbb{Q}(t)$ -linear combination of 1 and x . This can be done in such a way that the degree of the discriminant is minimal. We can read out some properties of the surface directly from the Weierstrass equation. For instance, if $p_a(X)$ is the arithmetic genus of X , then $p_a(X) + 1 = \chi(\mathcal{O}_X)$ is the minimal n such that $\deg a_i \leq ni$ for $i = 1, 2, 3, 4, 6$. In particular, for a K3 surface X , we need to have $\deg a_i \leq 2i$ (the case $n = 1$ corresponds to a rational elliptic surface).

All but finitely many of the fibers of the elliptic surface are nonsingular and hence elliptic curves. Tate's algorithm [T] allows us to compute the description of the singular fibers.

Each fiber is one of the types shown in the figure below. We note that the reducible fibers are unions of nonsingular rational curves, and they occur in configurations as shown below. The dual graph of the components is an extended Dynkin diagram of type A , D or E . The lattices labeled below are the ones generated by the non-identity components of the fiber in $NS(X)$. The subscript in the root lattices indicated below is the rank of the lattice, which is also the number of non-identity components. Thus, the I_n fiber has n components, whereas the I_n^* fiber has $n + 5$ components.



The Néron-Severi lattice of X is generated by the classes of all the sections of π (i.e. the Mordell-Weil group of X) considered as curves on the surface X , together with the class F of a fiber, and all the non-identity components of the reducible fibers. Let $R = \{v \in C(\mathbb{C}) \mid F_v \text{ is reducible}\}$, and for each $v \in R$, let $F_v = \pi^{-1}(v) = \Theta_{v,0} + \sum_{i=1}^{m_v-1} \mu_{v,i} \Theta_{v,i}$, where $\Theta_{v,0}$ is the component which intersects the identity, and the other $\Theta_{v,i}$ are the non-identity components. Let O be the zero section. The intersection pairing satisfies:

- (1) for any section P , $P^2 = O^2 = -\chi$,
- (2) $P \cdot F = O \cdot F = 1$,
- (3) $F^2 = 0$,
- (4) $O \cdot \Theta_{v,i} = 0$ for $i \geq 1$,
- (5) $\Theta_{v,i} \cdot \Theta_{w,j} = 0$ for $v \neq w$.

The intersection pairing for $\Theta_{v,i}$ and $\Theta_{v,j}$ is -2 if $i = j$, and $0, 1, 2$ if $i \neq j$ according to the figures above (2 occurs only for types I_2 and III). For a general section P , the intersection pairing with each $\Theta_{v,i}$ can be computed locally. In particular, for each v exactly one of the intersection numbers is 1, for some i such that $\mu_{v,i} = 1$, and the others vanish. The rank of the Néron-Severi group is given by the formula

$$\rho = r + 2 + \sum_v m_v - 1.$$

The discriminant of the sublattice T generated by the non-identity components of all the fibers is $\prod_v m_v^{(1)}$, where $m_v^{(1)}$ is the number of multiplicity one components of F_v .

4. GENUS 2 CURVES

4.1. Background: Moduli and invariants. Here we describe the basic geometry and moduli of curves of genus 2. For more background we refer the reader to [CF], [Cl], [I], [Me]. Let C be such a curve defined over a field k of characteristic zero. Then the canonical K_C bundle of C has degree 2 and $h^0(C, K_C) = 2$. That is, the corresponding complete linear system is a g_2^1 (and it is the unique g_2^1). We therefore have a map

$$x : C \rightarrow \mathbb{P}^1$$

which is ramified at 6 points by the Riemann-Hurwitz formula, and the function field of C is a quadratic extension of $k(x)$. Therefore, we may write the equation of C as

$$y^2 = f(x) = \sum_{i=0}^6 f_i x^i.$$

The roots of the sextic are the six ramification points of the map $C \rightarrow \mathbb{P}^1$. Their pre-images on C are the six Weierstrass points.

Now, the isomorphism class of C over $\bar{k}$, the algebraic closure of k , is determined by the isomorphism class of the sextic $f(x)$, where two sextics are equivalent if there is a transformation in $PGL_2(\bar{k})$ which takes the set of roots (considered inside $\mathbb{P}^1$) to the roots of the other.

Clebsch was the first to determine the invariants of binary sextics. He defined invariants of I_2, I_4, I_6, I_{10} of weights 2, 4, 6, 10 respectively. That is, I_d has degree d in the coefficients of f , and if f transforms to g under the action of $GL_2(\bar{k})$, then there is an element $r \in \bar{k}$ such that $I_d(g) = r^d I_d(f)$. Clebsch and Bolza showed that these invariants determined the sextic up to $\bar{k}$ -equivalence. Therefore, the point $(I_2(f) : I_4(f) : I_6(f) : I_{10}(f))$ in weighted projective space determines the isomorphism class of C . In fact, C and C' are isomorphic over k iff there is an $r \in k^*$ such that $I_d(f') = r^d I_d(f)$. Igusa generalized Clebsch's theory to hold in all characteristics by defining choosing a different algebraic equation for the curve C (through an embedding as a quartic in $\mathbb{P}^2$ with one node) and defining invariants J_2, J_4, J_6, J_8 and J_{10} . He thus obtained a moduli space of genus two curves defined over $\text{Spec } \mathbb{Z}$. The invariants $I_2, \dots, I_{10}$ are called the Igusa-Clebsch invariants whereas $J_2, \dots, J_{10}$ are called the Igusa invariants of C . We will use the former, since we are working over a field of characteristic zero.

Remark 12. If the Igusa-Clebsch invariants of a curve C lie in a field k , it does not necessarily mean that C can be defined over k : there is usually an obstruction in $Br_2(k)$. But C can always be defined over a quadratic extension of k .

4.2. Kummer surface. Let C be a curve of genus 2, which we can write as

$$y^2 = f(x) = \sum_{i=0}^6 f_i x^i$$

Let $\theta_i, i = 1, \dots, 6$ be the roots of the sextic, so that

$$f(x) = f_6 \prod_{i=1}^6 (x - \theta_i).$$

We shall concern ourselves with the embedding of the singular Kummer surface as a quartic in $\mathbb{P}^3$, which comes from the complete linear system 2Θ , twice the theta divisor which defines the principal polarization. We shall use the formulas from [CF]. The quartic is given by the equation

$$K(z_1, z_2, z_3, z_4) = K_2 z_4^2 + K_1 z_4 + K_0 = 0.$$

where

$$K_2 = z_2^2 - 4z_1 z_3,$$

$$K_1 = -4z_1^3 f_0 - 2z_1^2 z_2 f_1 - 4z_1^2 z_3 f_2 - 2z_1 z_2 z_3 f_3 - 4z_1 z_3^2 f_4 - 2z_2 z_3^2 f_5 - 4z_3^3 f_6,$$

$$\begin{aligned} K_0 = & -4z_1^4 f_0 f_2 + z_1^4 f_1^2 - 4z_1^3 z_2 f_0 f_3 - 2z_1^3 z_3 f_1 f_3 - 4z_1^2 z_2^2 f_0 f_4 \\ & + 4z_1^2 z_2 z_3 f_0 f_5 - 4z_1^2 z_2 z_3 f_1 f_4 - 4z_1^2 z_3^2 f_0 f_6 + 2z_1^2 z_3^2 f_1 f_5 \\ & - 4z_1^2 z_3^2 f_2 f_4 + z_1^2 z_3^2 f_3^2 - 4z_1 z_2^2 f_0 f_5 + 8z_1 z_2^2 z_3 f_0 f_6 \\ & - 4z_1 z_2^2 z_3 f_1 f_5 + 4z_1 z_2 z_3^2 f_1 f_6 - 4z_1 z_2 z_3^2 f_2 f_5 - 2z_1 z_3^3 f_3 f_5 \\ & - 4z_2^4 f_0 f_6 - 4z_2^3 z_3 f_1 f_6 - 4z_2^2 z_3^2 f_2 f_6 - 4z_2 z_3^3 f_3 f_6 - 4z_3^4 f_4 f_6 + z_3^4 f_5^2. \end{aligned}$$

The 16 singular points define ordinary double points on the quartic, which are called nodes. These are given explicitly by the coordinates

$$\begin{aligned} p_0 &= (0 : 0 : 0 : 1) \\ p_{ij} &= (1 : \theta_i + \theta_j : \theta_i \theta_j : \beta_0(i, j)) \end{aligned}$$

for $1 \leq i < j \leq 6$.

Here $\beta_0(i, j)$ is defined as follows. Let

$$f(x) = (x - \theta_i)(x - \theta_j)h(x) \text{ with } h(x) = \sum_{n=0}^4 h_n x^n.$$

Then

$$\beta_0(i, j) = -h_0 - h_2(\theta_i \theta_j) - h_4(\theta_i \theta_j)^2.$$

The singular point p_0 comes from the 0 point of the Jacobian, whereas the p_{ij} comes from the 2-torsion point which is the difference of divisors $[(\theta_i, 0)] - [(\theta_j, 0)]$ corresponding to two distinct Weierstrass points on C . The sixteen singular points are called **nodes**.

There are also sixteen hyperplanes in $\mathbb{P}^3$ which are tangent to the Kummer quartic. These are called **tropes**. Each trope intersects the quartic in a conic with multiplicity 2, and contains 6 nodes. Conversely, each node is contained in exactly 6 tropes. This beautiful configuration is called the $(16, 6)$ Kummer configuration.

The explicit formulas for the tropes are as follows. Six of the tropes are given by

$$\theta_i^2 z_1 - \theta_i z_2 + z_3 = 0.$$

We call this trope T_i . It contains the nodes p_0 and p_{ij} . The remaining ten tropes are labeled T_{ijk} and correspond to partitions of $\{1, 2, 3, 4, 5, 6\}$ into two sets of three, say $\{i, j, k\}$ and its complement $\{l, m, n\}$. Set

$$G(X) = (x - \theta_i)(x - \theta_j)(x - \theta_k) = \sum_{r=0}^3 g_r x^r,$$

$$H(X) = (x - \theta_l)(x - \theta_m)(x - \theta_n) = \sum_{r=0}^3 h_r x^r.$$

Then the equation of T_{ijk} is

$$f_6(g_2 h_0 + g_0 h_2) z_1 + f_6(g_0 + h_0) z_2 + f_6(g_1 + h_1) z_3 + z_4 = 0.$$

The Néron-Severi lattice of the nonsingular Kummer contains classes of rational curves E_0 and E_{ij} coming from the nodes, and C_i and C_{ijk} coming from the tropes. We will denote the lattice generated by these as $\Lambda_{(16,6)}$. It has signature $(1, 16)$ and discriminant 2^6 and is the Néron-Severi lattice of the Kummer surface of a generic principally polarized abelian surface.

Let L be the class of a hyperplane section. Furthermore, since T_{ijk} only depends on the partition $\{1, 2, 3, 4, 5, 6\} = \{i, j, k\} \cup \{l, m, n\}$, we may assume $i = 1$ and set $C_{jk} = C_{1jk}$ for $1 \leq j < k \leq 6$. Also set $C_{1j} = C_j$ for $1 < j \leq 6$ and $C_0 = C_1$ to agree with the notation in [Na]. We have the following intersection numbers and relations in the Néron-Severi lattice.

$$\begin{aligned} L^2 &= 4, \\ E_0^2 &= -2, \\ E_{ij}^2 &= -2, \\ E_0 \cdot E_{ij} &= 0, \\ E_{ij} \cdot E_{kl} &= 0, \{i, j\} \neq \{k, l\}, \\ C_0 &= (L - E_0 - \sum_k E_{1k})/2, \\ C_{1j} &= (L - E_0 - \sum_{k \neq j} E_{jk})/2, \\ C_{jk} &= (L - E_{ij} - E_{jk} - E_{ik} - E_{lm} - E_{mn} - E_{ln})/2 \end{aligned}$$

where $\{l, m, n\}$ is the complementary set to $\{1, j, k\}$.

We consider the following construction outlined in [Na]. Projection to a hyperplane from p_0 defines a 2 to 1 map of the Kummer to $\mathbb{P}^2$, and thus identifies the Kummer as a double cover of $\mathbb{P}^2$, ramified along the union of six lines, which are the projections of the conics C_i (or the tropes T_i). The exchange of sheets gives an involution of the lattice, which acts by

$$\begin{aligned} E_0 &\mapsto 2L - 3E_0, \\ E_{ij} &\mapsto E_{ij}, \\ L &\mapsto 3L - 4E_0. \end{aligned}$$

We can explicitly write down the projection to $\mathbb{P}^2$ as $(x_1, x_2, x_3, x_4) \mapsto (x_1, x_2, x_3)$. The involution which is the exchange of sheets is $(x_1, x_2, x_3, x_4) \mapsto (x_1, x_2, x_3, -x_4)$. Let q_0, q_{ij} be the projections of the p_0, p_{ij} .

5. FAMILY OF ELLIPTIC K3 SURFACES ASSOCIATED TO GENUS 2 CURVES

5.1. Elliptic surfaces with E_8 and E_7 fibers. Let X be an elliptic K3 surface with bad fibers of type E_8 and E_7 at ∞ and 0 respectively. A generic such K3 surface has a Néron-Severi lattice $NS(X) \cong U \oplus E_8(-1) \oplus E_7(-1)$ by Shioda's explicit description of the Néron-Severi lattice of an elliptic surface. This lattice has rank 17, signature $(1, 16)$ and discriminant 2. The transcendental lattice T_X has rank 5, signature $(2, 3)$ and discriminant 2. We deduce that $T_X \cong U^2 \oplus \langle -2 \rangle$.

The transcendental lattice of a generic principally polarized abelian surface, that is, the Jacobian $J(C)$ for C a generic curve of genus 2, also satisfies the same property, since the Néron-Severi of $J(C)$ is generated by the theta divisor, which has self-intersection 2 by the genus formula on the abelian surface

$$2 = 2g - 2 = C \cdot (C + K) = C^2.$$

Therefore the orthogonal complement in $H^2(J(C), \mathbb{Z}) \cong U^3$ is exactly $U^2 \oplus \langle -2 \rangle$. We expect that the elliptic K3 surface X has a Shioda-Inose structure such that the quotient by the Nikulin involution gives the Kummer surface of a principally polarized abelian surface $\text{Km}(J(C))$. Galluzzi, Lombardo and Dolgachev [GLD] prove that, in fact, X corresponds to a unique C up to isomorphism. However, an explicit identification of the quotient as a Kummer surface was not known. Below, we give an explicit construction of the correspondence.

We begin with the K3 surface X given by the equation

$$y^2 = x^3 + t^3(at + a')x + t^5(b''t^2 + bt + b').$$

It is easily checked that the surface X has an II^* or E_8 fiber at $t = \infty$ and a III^* or E_7 fiber at $t = 0$. Now we describe another elliptic fibration on X .

Applying the transformation $(x, y, t) = (x't'^2/b''^2, y't'^2/b''^3, t'/b'')$, we get

$$y'^2 = t'^3 + (x'^3 + ax' + b)t'^2 + b''(a'x' + b')t'$$

and again replacing (x', y', t') by (x, y, t) for convenience gives finally

$$y^2 = t^3 + (x^3 + ax + b)t^2 + b''(a'x + b')t$$

which is an elliptic surface over the x -line with an I_{10}^* or D_{14} fiber at $x = \infty$, an I_2 or A_1 fiber at $x = -b'/a'$ and a 2-torsion section $(y, t) = (0, 0)$. The translation by the 2-torsion section is a Nikulin involution. We write down the isogenous elliptic surface Y as

$$y^2 = t^3 - 2(x^3 + ax + b)t^2 + ((x^3 + ax + b)^2 - 4b''(a'x + b'))t.$$

This is an elliptic surface over the x -line with an I_5^* or D_9 fiber at $x = \infty$ and I_2 or A_1 fibers at the roots of the sextic $(x^3 + ax + b)^2 - 4b''(a'x + b')$, and with a 2-torsion section $(t, y) = (0, 0)$. The Néron-Severi lattice of a generic such surface

has signature $(1, 16)$ and discriminant $4 \cdot 2^6/2^2 = 2^6$. In fact, we will identify it with the Néron-Severi lattice of a generic Kummer surface (which we call the $(16, 6)$ lattice) in a later section. This will lead to the identification of the Kummer surface of $J(C)$ as an elliptic K3 surface with a bad fiber of type I_5^* at ∞ , with I_2 fibers at the roots of a sextic derived from C , and with a 2-torsion section.

Remark 13. If E is an elliptic curve with a 2-torsion point, written in the Weierstrass form

$$y^2 = x^3 + ax^2 + bx$$

with a 2-torsion point $P = (0, 0)$, then the 2-isogenous curve $E' = E/\{O, P\}$ is given by

$$Y^2 = X^3 - 2aX^2 + (a^2 - 4b)X.$$

The isogeny $\phi : E \rightarrow E'$ is given by

$$(x, y) \rightarrow \left(\frac{y^2}{x^2}, \frac{y(b - x^2)}{x^2} \right).$$

The dual isogeny $\hat{\phi}$ is given by

$$(X, Y) \rightarrow \left(\frac{Y^2}{4X^2}, \frac{Y(a^2 - 4b - X^2)}{8X^2} \right).$$

5.2. Main theorem. In this section, we state the main theorem, which gives the parameters (a, a', b, b', b'') of the K3 elliptic surface with E_8 and E_7 fibers (so far conjecturally) associated to a genus 2 curve C in terms of the Igusa-Clebsch invariants of C . The proof will be given in the following sections.

Theorem 14. *Let C be a curve of genus two, and $Y = \text{Km}(J(C))$ the Kummer surface of its Jacobian. Let I_2, I_4, I_6, I_{10} be the Igusa-Clebsch invariants of Y . Then there is an elliptic fibration on Y for which the Weierstrass equation may be written*

$$y^2 = x^3 - 2 \left(t^3 - \frac{I_4}{12}t + \frac{I_2I_4 - 3I_6}{108} \right) x^2 + \left(\left(t^3 - \frac{I_4}{12}t + \frac{I_2I_4 - 3I_6}{108} \right)^2 + I_{10} \left(t - \frac{I_2}{24} \right) \right) x.$$

There is an elliptic K3 surface X given by

$$y^2 = x^3 - t^3 \left(\frac{I_4}{12}t + 1 \right) x + t^5 \left(\frac{I_{10}}{4}t^2 + \frac{I_2I_4 - 3I_6}{108}t + \frac{I_2}{24} \right)$$

with fibers of type E_8 and E_7 at $t = \infty$ and $t = 0$ respectively, and a Nikulin involution on X , such that the quotient K3 surface is Y .


Remark 15. Note that the correspondence of the K3 surface X with the genus 2 curve C is Galois invariant, i.e. X is defined over the field of definition of C . The proof of the theorem will involve making some non-Galois invariant choices (for the level 2 structure of the Jacobian of C) but we will show that X as well as the Shioda-Inose structure on X is independent of these choices.

Remark 16. The Nikulin involution on X may be written as follows:

$$(x, y, t) \mapsto \left(\frac{16x(-x + I_2t^2/24)^2}{I_{10}^2 t^8}, \frac{-64y(-x + I_2t^2/24)^3}{I_{10}^3 t^{12}}, \frac{4(-x + I_2t^2/24)}{I_{10} t^3} \right).$$


5.3. Néron-Severi lattices. In this and the next section, we give the details of how to put an elliptic fibration on the Kummer surface of a Jacobian of a curve of genus 2, with a 2-torsion section, a I_5^* fiber and six I_2 fibers. We first identify the Néron-Severi lattices involved (namely $\Lambda_{(16,6)}$ and $(D_9 \oplus A_1^6 \oplus U)^+$). We use the construction of [Na], which gives an embedding of the lattice $N \oplus E_8(-1)$ inside $\Lambda_{(16,6)}$.

First, we start with the Néron-Severi lattice of the K3 surface X which has E_8 and E_7 fibers. The roots of the $NS(X)$ which correspond to the smooth rational curves on X are drawn below (we use the notation from [GLD]). These are in fact all the roots corresponding to smooth rational curves on X , by a result of Nikulin [N2].



There is an elliptic fibration on X which has $R_8 + 2R_7 + 3R_6 + 4R_5 + 5R_4 + 6R_3 + 4R_2 + 2R_1 + 3R_0$ as an II^* or E_8 fiber, $N_7 + 2N_6 + 3N_5 + 4N_4 + 3N_3 + 2N_2 + N_1 + 2N_0$ as a III^* or E_7 fiber, and S as the zero section. This is the fibration over $\mathbb{P}_t^1$. The fibration over $\mathbb{P}_x^1$ has the I_{10}^* or D_{14} fiber given by $R_0 + R_2 + 2(R_3 + R_4 + R_5 + R_6 + R_7 + R_8 + S + N_1 + N_2 + N_3 + N_4) + N_0 + N_5$, an I_2 or A_1 fiber $A + N_7$, a 2-torsion section (say R_1) and a zero section N_6 .


The Nikulin involution σ is translation by the 2-torsion section. It reflects the above picture about its vertical axis of symmetry. There are two obvious copies of $E_8(-1)$ switched by σ , namely the sublattices of $NS(X)$ generated by the roots $\{S, N_1, N_2, N_3, N_4, N_0, N_5, N_6\}$ and $\{R_7, R_6, R_5, R_4, R_3, R_0, R_2, R_1\}$. Next, we write down some roots on $NS(Y)$, where Y is the quotient K3 surface of X by the involution. As we have described, Y has six I_2 or A_1 fibers $Q_{13} + Q_{14}, \dots, Q_{23} + Q_{24}$, a I_5^* or D_9 fiber, namely $Q_1 + Q_2 + 2(Q_3 + Q_4 + Q_5 + Q_6 + Q_7 + Q_8) + Q_9 + Q_{10}$, with zero section $O = Q_{11}$ and a 2-torsion section $T = Q_{12}$.



It is easily checked that the rational components of the E_8 fiber described above map as following: $N_6 \mapsto O$ (recall that N_6 is the zero section of the D_{14} fibration, on which the quotient map is an isogeny of elliptic surfaces), $N_5 \mapsto Q_9$, $N_4 \mapsto Q_8$, $N_0 \mapsto Q_{10}$, $N_3 \mapsto Q_7$, $N_2 \mapsto Q_6$, $N_1 \mapsto Q_5$, $S \mapsto Q_4$. Hence, we see a natural copy of E_8 within the Néron-Severi lattice of Y . On the other hand, we can also see eight roots orthogonal to all the generators of E_8 as well as to each other, namely $Q_{14}, Q_{16}, Q_{18}, Q_{20}, Q_{22}, Q_{24}, Q_1$ and Q_2 .

Now, we use the construction of Naruki [Na] which gives an explicit embedding of $N \oplus E_8(-1)$ inside the Néron-Severi lattice of a Kummer surface of a generic principally polarized abelian surface, or $\Lambda_{(16,6)}$. We extend this construction to get an identification of $\Lambda_{(16,6)}$ with $NS(Y)$, i.e. the lattice generated by the roots in the diagram above.

The identification is as follows:



Here $\alpha(C_{23}) = C_{23} + L - 2E_0$.

The class of the fiber is

$$\begin{aligned} F &= C_{23} + \alpha(C_{23}) + 2(E_{23} + C_{12} + E_{26} + C_{16} + E_{16} + C_0) + E_{15} + E_{14} \\ &= 5(L - E_0) - 3E_{12} - 2(E_{13} + E_{46} + E_{56}) - (E_{24} + E_{25} + E_{36} + E_{45}) \end{aligned}$$

and $e_1, \dots, e_6, f_1, \dots, f_6$ are given by

$$\begin{aligned} e_1 &= (L - E_0) - (E_{12} + E_{46}) \\ e_2 &= 2(L - E_0) - (E_{12} + E_{13} + E_{24} + E_{46} + E_{56}) \\ e_3 &= 3(L - E_0) - 2E_{12} - (E_{13} + E_{24} + E_{36} + E_{45} + E_{46} + E_{56}) \\ e_4 &= 4(L - E_0) - 2(E_{12} + E_{13} + E_{46}) - (E_{24} + E_{25} + E_{36} + E_{45} + E_{56}) \\ e_5 &= 5(L - E_0) - 3E_{12} - 2(E_{13} + E_{46} + E_{56}) - (E_{24} + E_{25} + E_{34} + E_{36} + E_{45}) \\ e_6 &= E_{35} \\ (f_i &= F - e_i \text{ for all } i) \\ f_4 &= (L - E_0) - (E_{12} + E_{56}) \\ f_3 &= 2(L - E_0) - (E_{12} + E_{13} + E_{25} + E_{46} + E_{56}) \\ f_2 &= 3(L - E_0) - 2E_{12} - (E_{13} + E_{25} + E_{36} + E_{45} + E_{46} + E_{56}) \\ f_1 &= 4(L - E_0) - 2(E_{12} + E_{13} + E_{56}) - (E_{24} + E_{25} + E_{36} + E_{45} + E_{46}) \\ f_6 &= 5(L - E_0) - 3E_{12} - 2(E_{13} + E_{46} + E_{56}) - (E_{24} + E_{25} + E_{34} + E_{36} + E_{45}) \\ f_5 &= E_{34}. \end{aligned}$$

Notice that under the simple transposition (45) of indices we have the permutation of fibers $\tau = (14)(23)(56)$ and in fact $e_i \mapsto f_{\tau(i)}$, $f_i \mapsto e_{\tau(i)}$.

5.4. Completion of proof. Next, we describe how to use all this information from the Néron-Severi group to construct x , y and t in the Weierstrass equation for $Y = \text{Km}(J(C))$

$$y^2 = x^3 + a(t)x^2 + b(t)x$$

Consider the class of the fiber $F \in NS(\text{Km}(J(C)))$, given by

$$F = 5(L - E_0) - 3E_{12} - 2(E_{13} + E_{46} + E_{56}) - (E_{24} + E_{25} + E_{36} + E_{45}).$$

We can write down the parameter on the base by computing explicitly the sections of $H^0(Y, \mathcal{O}_Y(F))$. This linear system consists of (the pullback of) quintics passing through the points q_0 and q_{ij} which pass through $q_{24}, q_{25}, q_{36}, q_{45}$, having a double point at q_{13}, q_{46}, q_{56} and a triple point at q_{12} . This linear system is 2-dimensional, and taking the ratio of two linearly independent sections gives us the parameter t on the base, $\mathbb{P}^1$, for the elliptic fibration. Now, t is only determined up to the action of PGL_2 , but the first restriction we make is to put the I_5^* fiber at $t = \infty$, which fixes t up to affine linear transformations. Any elliptic K3 surface with a 2-torsion section can be written in the form

$$y^2 = x^3 - 2q(t)x^2 + p(t)x$$

with $p(t)$ of degree at most 8 and $q(t)$ of degree at most 4. The 2-torsion section is $(x, y) = (0, 0)$. The discriminant of this elliptic surface is a multiple of $p^2(q^2 - p)$. In fact, we see that p must have degree exactly 6, because there are six I_2 fibers that the zero and 2-torsion sections meet in different components. The positions $t_1, \dots, t_6$ of the I_2 fibers are the roots of the polynomial $p(t) = p_0 \prod_{i=1}^6 (x - t_i)$. Now t is determined up to transformations of the form $t \mapsto at + b$. To have exactly

a I_5^* fiber at ∞ , we must have $p(t) = q(t)^2 + r(t)$ where $q(t)$ is a monic cubic polynomial and $r(t)$ is a *linear* polynomial in t . We can further fix t up to scalings $t \mapsto at$ by translating t so that the quadratic term of $q(t)$ vanishes. We notice that the top coefficient p_0 of $p(t)$ is a square, and so by scaling t, x, y appropriately, we may assume $p_0 = 1$, i.e. that $p(t)$ and $q(t)$ are monic.

Now we describe how to obtain x . It is a Weil function, so that the horizontal component of its divisor equals $2T - 2O$, and the vertical component is uniquely determined by that fact that (x) is linearly (and hence numerically) equivalent to zero. So we deduce that the divisor of x is $2T - 2O + Q_{10} - Q_9 + Q_{14} + Q_{16} + Q_{18} + Q_{20} + Q_{22} + Q_{24} - 3F_0$, where

$$F_0 = Q_1 + Q_2 + 2(Q_3 + Q_4 + Q_5 + Q_6 + Q_7 + Q_8) + Q_9 + Q_{10}$$

is the D_9 fiber.

To convert this to formulas, we compute the functions which cut out $Q_{16}, \dots, Q_{24}$, F_0 , T and O . There is a quintic s_1 which cuts out $O = C_{14}$. Now, notice that the D_9 fiber contains C_{12}, C_{16} and C_0 . Therefore s_1 is divisible by T_2, T_6 and T_1 . We write

$$s_1 = q_1 T_1 T_2 T_6$$

with a quadratic q_1 . Next, we know that T_4 cuts out $C_{14} = O$ and T_5 cuts out $C_{15} = T$. To find, for instance, the function which cuts out e_2 , we find the quadratic (unique up to constants) which passes through $q_{12}, q_{13}, q_{24}, q_{46}, q_{56}$. Call this function e_2 , by abuse of notation. Similarly, we find $e_1, \dots, e_5$. We also note that the factor of T_5 in the numerator of x , which gives a zero along T , also gives a zero along $e_6 = E_{35}$ owing to the fact that $T = C_{15}$ intersects E_{35} (recall that we are working with the singular Kummer surface, on which the image of the curve E_{35} is just a single point).

Putting everything together, we can write x up to scaling as a quotient of two homogeneous polynomials of degree 16 as follows:

$$x = \frac{e_1 e_2 e_3 e_4 e_5 T_5}{s_1^3 T_4} = \frac{e_1 e_2 e_3 e_4 e_5 T_5}{(T_1 T_2 T_6 q_1)^3 T_4}$$

Finally, we have to scale x and t so that $x^3 + a(t)x^2 + b(t)x$ becomes a square of a function y on the Kummer.

We note that in the equation of the Kummer

$$K_2 z_4^2 + K_1 z_4 + K_0 = 0$$

we can complete the square for z_4 to obtain

$$(K_2 z_4 + K_1/2)^2 = K_1^2/4 - K_0 K_2 = 4T_1 T_2 T_3 T_4 T_5 T_6$$

We let y be a constant multiple of

$$\frac{e_1 e_2 e_3 e_4 e_5 (K_2 z_4 + K_1/2)}{T_1^5 T_2^3 T_4^2 T_6^4 q_1^2},$$

a quotient of two homogenous polynomials of degree 18, and verify that this makes the Weierstrass equation hold. The computation is carried out in a Maxima program, which is available from the arXiv.org e-print archive. This paper is available as `math.AG/0701669`. To access the auxiliary file download the source file for the paper. That will produce not only the $\text{\LaTeX}$ files for the paper but also the computer algebra code. The code takes about half an hour to run on a 2.1 GHz computer.

We noted earlier that the permutation (45) on the A_1 fibers by $\tau = (14)(23)(56)$, and takes e_i to $f_{\tau(i)}$. That is, it switches the components intersecting the identity and 2-torsion sections as well. In addition, it switches the zero section C_{14} and the 2-torsion section C_{15} , and on the D_9 fibers it switches the two near leaves E_{15} and E_{14} , namely, again the components intersecting T and O . On the other hand, consider the action on $NS(Y)$ induced by the translation by T . Under this map, T and O get swapped, the 2-torsion and identity components of the D_9 and A_1 fibers all get switched, and the far leaves of the D_9 fiber also get switched (this can be seen, for instance, from the fact that the simple components of the special D_9 fiber form a group compatible with the group law on the generic elliptic curve). The locations of the A_1 fibers themselves are fixed. Therefore the effect of the permutation 45 is the same as translation by 2-torsion composed with a pure involution (14)(23)(56) of the A_1 fibers and a switch of the far leaves of the D_9 fiber. Since the far leaves of the D_9 fiber are switched by the Galois involution that multiplies the square root of $b'' = I_{10}/4$ by -1 , this tells us that we have the correct quadratic twist, since I_{10} is within a square factor of the discriminant of the sextic. That is, making a different choice of 2-level structure while describing the fibration on the Kummer surface Y would have given us the same answer for X !

6. AN APPLICATION TO THE MODULI OF SIX POINTS IN $\mathbb{P}^1$

The construction above gave us a correspondence of sextics

$$f(x) = \sum f_i x^i = \prod (x - x_i)$$

and

$$g(x) = \left(x^3 - \frac{I_4}{12}x + \frac{I_2 I_4 - 3I_6}{108} \right)^2 + I_{10} \left(x - \frac{I_2}{24} \right).$$

Therefore, over an algebraically closed field, we get a birational map from the moduli space of 6 points in $\mathbb{P}^1$ (i.e. the quotient of $(\mathbb{P}^1)^6$ under the action of PGL_2 and S_6) with the space of roots up to scaling of

$$(x^3 + ax + b)^2 + (a'x + b')$$

as a, b, a', b' vary (we suppressed b'' since it just scales a' and b'). This latter space is cut out inside $\mathbb{P}^5 = \{(X_1 : X_2 : X_3 : X_4 : X_5 : X_6)\}$ by the hyperplane $\sigma_1(X) = X_1 + \dots + X_6 = 0$ and the quartic hypersurface $\sigma_2(X)^2 = 4\sigma_4(X)$, where σ_2 and σ_4 are the second and fourth elementary symmetric functions of the X_i . Thus, we get a model as a singular quartic threefold in $\mathbb{P}^4$, which is known in the literature as the Igusa quartic. Here the Igusa quartic arises naturally in the context of Shioda-Inose structures on K3 surfaces.

There is no simple one-one correspondence between the roots of $f(x)$ and $g(x)$, since the two actions of S_6 acting by the permutation representation on the six roots of $f(x)$ on the six roots of $g(x)$ are related by an outer automorphism. To see this, we recall from the last section that the permutation (45) on the roots of $f(x)$ (the Weierstrass points) acts on the roots of $g(x)$ (which are the locations of the A_1 fibers) by the permutation (14)(23)(56). By symmetry, all the transpositions of S_6 act by a product of three transpositions on the roots of $g(x)$. Thus we get a homomorphism $S_6(f) \rightarrow S_6(g)$ which is an outer automorphism.

ACKNOWLEDGEMENTS

I thank Noam Elkies and Barry Mazur for suggesting the study of K3 surfaces as a thesis topic and for many insightful remarks and helpful conversations. I also thank Tetsuji Shioda for discussions and correspondence on elliptic surfaces, and Igor Dolgachev for helpful remarks and pointing out some useful references. I also thank Charles Doran for bringing his work with Adrian Clinger to my attention.

REFERENCES

- [BHPV] W. P. Barth, K. Hulek, C. A. M. Peters and A. Van de Ven, *Compact complex surfaces*, Second edition, Springer-Verlag, Berlin, 2004.
- [CF] J. W. F. Cassels and E. V. Flynn, *Prolegomena to a middlebrow arithmetic of curves of genus 2*, London Mathematical Society Lecture Note Series, 230. Cambridge University Press, Cambridge, 1996.
- [Cl] A. Clebsch, *Zur Theorie der binären algebraischen Formen*, Verlag von B. G. Teubner, Leipzig, 1872.
- [CD] A. Clinger and C. Doran, *Modular invariants for lattice polarized K3 surfaces*, preprint, 2006.
- [E] N. D. Elkies, *On a family of K3 surfaces of rank at least 18*, preprint, 2005.
- [GLD] F. Galluzzi and G. Lombardo, *Correspondences between K3 surfaces. With an appendix by Igor Dolgachev*, Michigan Math. J. **52** (2004), no. 2, 267–277.
- [I] J. Igusa, *Arithmetic variety of moduli for genus two*, Ann. of Math. (2) **72** (1960), 612–649.
- [In] H. Inose, *Defining equations of singular K3 surfaces and a notion of isogeny*, Proceedings of the International Symposium on Algebraic Geometry (Kyoto Univ., 1977), 495–502.
- [IS] H. Inose and T. Shioda, *On singular K3 surfaces*, Complex Analysis and algebraic geometry 75–81, Iwanami Shoten, Tokyo, 1977.
- [K] *Géométrie des surfaces K3: modules et périodes. (French) [Geometry of K3 surfaces: moduli and periods]*, Papers from the seminar held in Palaiseau, October 1981–January 1982. Astérisque No. 126 (1985). Société Mathématique de France, Paris, 1985. pp. 1–193.
- [MS] N. Mitani and T. Shioda, *Singular abelian surfaces and binary quadratic forms*, Classification of algebraic varieties and compact complex manifolds, pp. 259–287. Lecture Notes in Math. Vol. 412, Springer, Berlin, 1974.
- [M1] D. R. Morrison, *On K3 surfaces with large Picard number*, Invent. Math. **75** (1984) no. 1, 105–121.
- [M2] D. R. Morrison, *Notes on the geometry of K3 surfaces*, Lectures delivered at the Scuola Matematica Interuniversitaria, Cortona, 1988.

- [Me] J.-F. Mestre, *Construction de courbes de genre 2 à partir de leurs modules*, Effective methods in algebraic geometry (Castiglione, 1990), 313–334, Progr. Math. 94, Birkhäuser Boston, Boston, MA, 1991.
- [Mi] J. Milnor, *On simply connected 4-manifolds*, Symposium internacional de topología algebraica (International symposium on algebraic topology), Universidad Nacional Autónoma de México and UNESCO, Mexico (1958) pp. 122–128
- [N1] V. Nikulin, *Finite groups of automorphisms of Kählerian surfaces of type K3*, Trudy Mosk. Math. Ob. **38** (1979), 75–137; Trans. Moscow Math. Soc. **38** (1980), 71–135.
- [N2] V. V. Nikulin, *Quotient-groups of groups of automorphisms of hyperbolic forms by subgroups generated by 2-reflections. Algebro-geometric applications*, Current problems in mathematics, v. 18, pp. 3–114, Akad. Nauk SSSR, Vsesoyuz. Inst. Nauchn. i Tekhn. Informatsii, Moscow, 1981; J. Soviet. Math. **22** (1983), 1401–1476.
- [Na] I. Naruki, *On metamorphosis of Kummer surfaces*, Hokkaido Math. J. **20** (1991), no. 2, 407–415.
- [S] T. Shioda, *On the Mordell-Weil lattices*, Comment. Math. Univ. St. Paul. **39** (1990), no. 2, 211–240.
- [Si] J. H. Silverman, *Advanced topics in the arithmetic of elliptic curves*, Graduate Texts in Mathematics, 151. Springer-Verlag, New York, 1994.
- [T] J. Tate, *Algorithm for determining the type of a singular fiber in an elliptic pencil*, Modular functions of one variable, IV (Proc. Internat. Summer School, Univ. Antwerp, Antwerp, 1972), pp. 33–52. Lecture Notes in Math. Vol. 476, Springer, Berlin, 1975.

MICROSOFT RESEARCH, ONE MICROSOFT WAY, REDMOND, WA 98052-6399

E-mail address: abhinavk@microsoft.com